

201-16124B

RECEIVED
CPPT CHIC

2005 DEC 34 AM 7:59

**APPENDIX I
ROBUST SUMMARIES**

I U C L I D

Data Set

Existing Chemical	: ID: 101144
Producer related part	
Company	: Epona Associates, LLC
Creation date	: 19.12.2005
Substance related part	
Company	: Epona Associates, LLC
Creation date	: 19.12.2005
Status	:
Memo	: MOCA
Printing date	: 22.12.2005
Revision date	:
Date of last update	: 22.12.2005
Number of pages	: 19
Chapter (profile)	: Chapter: 2.1, 2.2, 2.4, 2.5, 2.6.1, 3.1.1, 3.1.2, 3.3.1, 3.5, 4.1, 4.2, 4.3, 5.1.1, 5.1.2, 5.1.3, 5.1.4, 5.4, 5.5, 5.6, 5.8.1, 5.8.2
Reliability (profile)	: Reliability: without reliability, 1, 2, 3, 4
Flags (profile)	: Flags: without flag, confidential, non confidential, WGK (DE), TA-Luft (DE), Material Safety Dataset, Risk Assessment, Directive 67/548/EEC, SIDS

2. Physico-Chemical Data

Id 101144

Date 22.12.2005

2.1 MELTING POINT

Value : = 102 - 107 °C
Sublimation :
Method : other
Year : 2000
GLP : no
Test substance : as prescribed by 1.1 - 1.4

Reliability : (2) valid with restrictions
Data are taken from a secondary literature source (handbook)

Flag : Critical study for SIDS endpoint
19.12.2005 (2)

Value : = 110 °C
Sublimation :
Method : other
Year : 1983
GLP : no
Test substance : as prescribed by 1.1 - 1.4

Reliability : (1) valid without restriction
Data were obtained by modeling, SRC recommended value

19.12.2005 (56)

2.2 BOILING POINT

Value : = 378.9 °C at
Decomposition :
Method : other: calculation
Year : 1988
GLP : no
Test substance : as prescribed by 1.1 - 1.4

Reliability : (2) valid with restrictions
Data were obtained by modeling, estimated by PCCHEM-PCGEMS, SRC recommended value

Flag : Critical study for SIDS endpoint
19.12.2005 (51)

2.4 VAPOUR PRESSURE

Value : = .0000133 hPa at 25 °C
Decomposition :
Method :
Year : 1983
GLP : no data
Test substance : as prescribed by 1.1 - 1.4

Result : Vapor pressure = 1.0×10^{-5} mm Hg at 25 deg C

Reliability : (2) valid with restrictions
Source is peer-reviewed published data

Flag : Critical study for SIDS endpoint
21.12.2005 (50)

Value : < 0 hPa at 25 °C

2. Physico-Chemical Data

Id 101144

Date 22.12.2005

Decomposition	:		
Method	:	other (calculated)	
Year	:	1988	
GLP	:	no	
Test substance	:	as prescribed by 1.1 - 1.4	
Result	:	Vapor pressure = 1.32E-8 mm Hg at 25C Calc	
Reliability	:	(2) valid with restrictions Data were obtained by modeling, estimated by PCCHEM-PCGEMS, SRC recommended value	
19.12.2005			(51)
Value	:	< 0 hPa at 25 °C	
Decomposition	:		
Method	:	other (calculated)	
Year	:	2003	
GLP	:	no	
Test substance	:	as prescribed by 1.1 - 1.4	
Result	:	Vapor pressure = 2.86X10-7 mm Hg @ 25 deg C Calc	
Reliability	:	(2) valid with restrictions Data were obtained by modeling	
19.12.2005			(59)
Value	:	= .0000173 hPa at 60 °C	
Decomposition	:		
Method	:		
Year	:	1981	
GLP	:	no data	
Test substance	:	as prescribed by 1.1 - 1.4	
Result	:	Vapor pressure = 1.3 x 10E-5 mm Hg at 60 deg C	
Reliability	:	(2) valid with restrictions Source is peer-reviewed published data	
21.12.2005			(43)
Value	:	= .00173 hPa at 60 °C	
Decomposition	:		
Method	:	other (measured)	
Year	:	2001	
GLP	:	no	
Test substance	:	as prescribed by 1.1 - 1.4	
Result	:	Vapor pressure = 1.3X10-3 torr @ 60 deg C	
Reliability	:	(2) valid with restrictions Source is peer-reviewed published data	
21.12.2005			(1)
Value	:	= .0000466 hPa at 100 °C	
Decomposition	:		
Method	:		
Year	:	1983	
GLP	:	no data	
Test substance	:	as prescribed by 1.1 - 1.4	
Result	:	Vapor pressure = 3.5 x 10E-5 mm Hg at 100 deg C	
Reliability	:	(2) valid with restrictions Source is peer-reviewed published data	
20.12.2005			(50)
Value	:	= .0000719 hPa at 120 °C	
Decomposition	:		

2. Physico-Chemical Data

Id 101144

Date 22.12.2005

Method :
Year : 1981
GLP : no data
Test substance : as prescribed by 1.1 - 1.4

Result : Vapor pressure = 5.4×10^{-5} mm Hg at 120 deg C
Reliability : (2) valid with restrictions
Source is peer-reviewed published data

20.12.2005 (43)

2.5 PARTITION COEFFICIENT

Partition coefficient : octanol-water
Log pow : = 3.61 - 3.9 at °C
pH value : -
Method : other (calculated)
Year : 1978
GLP : no
Test substance : as prescribed by 1.1 - 1.4

Reliability : (2) valid with restrictions
EPA source document
Flag : Critical study for SIDS endpoint
19.12.2005 (27)

Partition coefficient : octanol-water
Log pow : = 3.94 at °C
pH value :
Method : other (calculated)
Year : 1988
GLP : no
Test substance : as prescribed by 1.1 - 1.4

Reliability : (2) valid with restrictions
Data were obtained by modeling, estimated by PCCHEM-PCGEMS, SRC
recommended value
19.12.2005 (51)

2.6.1 SOLUBILITY IN DIFFERENT MEDIA

Solubility in : Water
Value : = 13.9 mg/l at 24 °C
pH value :
concentration : at °C
Temperature effects :
Examine different pol. :
pKa : at 25 °C
Description :
Stable :
Deg. product :
Method : other
Year : 1986
GLP : no
Test substance : as prescribed by 1.1 - 1.4

Reliability : (2) valid with restrictions
Source is peer-reviewed published data
Flag : Critical study for SIDS endpoint

2. Physico-Chemical Data

Id 101144

Date 22.12.2005

19.12.2005

(61)

Solubility in : Water
Value : = 8.684 mg/l at 25 °C
pH value :
concentration : at °C
Temperature effects :
Examine different pol. :
pKa : at 25 °C
Description :
Stable :
Deg. product :
Method : other
Year : 2005
GLP : no
Test substance : as prescribed by 1.1 - 1.4

Result : Water Solubility Estimate from Log Kow (WSKOW v1.41):
Water Solubility at 25 deg C (mg/L): 8.684
log Kow used: 3.91 (expkow database)
no-melting pt equation used

Reliability : (2) valid with restrictions
Data were obtained by modeling

20.12.2005

(59)

3. Environmental Fate and Pathways

Id 101144

Date 22.12.2005

3.1.1 PHOTODEGRADATION

Type : air
Light source :
Light spectrum : nm
Relative intensity : based on intensity of sunlight
INDIRECT PHOTOLYSIS
Sensitizer : OH
Conc. of sensitizer :
Rate constant : = .0000000000775166 cm³/(molecule*sec)
Degradation : = 50 % after .1 day(s)
Deg. product :
Method : other (calculated)
Year : 2005
GLP : no
Test substance : as prescribed by 1.1 - 1.4

Result : Atmospheric Oxidation (25 deg C) [AopWin v1.91]:
Hydroxyl Radicals Reaction: OVERALL OH Rate Constant = 77.5166 E-12 cm³/molecule-sec

Half-Life = 0.138 Days (12-hr day; 1.5E6 OH/cm³)
Half-Life = 1.656 Hrs

Ozone Reaction: No Ozone Reaction Estimation

Reliability : (2) valid with restrictions
Data were obtained by modeling

Flag : Critical study for SIDS endpoint
21.12.2005 (59)

Type : air
Light source :
Light spectrum : nm
Relative intensity : based on intensity of sunlight
INDIRECT PHOTOLYSIS
Sensitizer : OH
Conc. of sensitizer : 500000 molecule/cm³
Rate constant : = .0000000000775 cm³/(molecule*sec)
Degradation : = 50 % after 5 hour(s)
Deg. product :
Method : other (calculated)
Year : 1993
GLP : no
Test substance : as prescribed by 1.1 - 1.4

Result : t1/2 = 5 hours ; rate constant = 7.75X10-11 cu cm/molecule-sec at 25 deg C

Reliability : (2) valid with restrictions
Data were obtained by modeling
20.12.2005 (38)

3.1.2 STABILITY IN WATER

Type : abiotic
t1/2 pH4 : at °C
t1/2 pH7 : > 1 year at 25 °C
t1/2 pH9 : at °C

3. Environmental Fate and Pathways

Id 101144
Date 22.12.2005

Deg. product :
Method : other: measurement
Year : 1988
GLP : no
Test substance : as prescribed by 1.1 - 1.4

Reliability : (2) valid with restrictions
EPA source document
Flag : Critical study for SIDS endpoint
19.12.2005

(17)

Type : abiotic
t1/2 pH4 : at °C
t1/2 pH7 : > 800 year at 25 °C
t1/2 pH9 : at °C
Deg. product :
Method : other
Year : 1989
GLP : no data
Test substance : as prescribed by 1.1 - 1.4

Result : 4,4'-Methylenebis(2-chloroaniline) is not expected to undergo hydrolysis based on a hydrolysis half-life of greater than 800 years at pH 7 and 25 deg C.
Reliability : (2) valid with restrictions
EPA source document
19.12.2005

(16)

3.3.1 TRANSPORT BETWEEN ENVIRONMENTAL COMPARTMENTS

Type : fugacity model level III
Media :
Air : % (Fugacity Model Level I)
Water : % (Fugacity Model Level I)
Soil : % (Fugacity Model Level I)
Biota : % (Fugacity Model Level II/III)
Soil : % (Fugacity Model Level II/III)
Method : other
Year : 2005

Result : Level III Fugacity Model:

	Mass Amount (percent)	Half-Life (hr)	Emissions (kg/hr)
Air	9.87e-005	3.31	1000
Water	15.5	1.44e+003	1000
Soil	82.1	1.44e+003	1000
Sediment	2.42	5.76e+003	0

Persistence Time: 1.59e+003 hr

Reliability : (2) valid with restrictions
Data were obtained by modeling
Flag : Critical study for SIDS endpoint
21.12.2005

(59)

3.5 BIODEGRADATION

Type : aerobic
Inoculum : activated sludge

3. Environmental Fate and Pathways

Id 101144

Date 22.12.2005

Concentration : 100 mg/l related to Test substance related to

Contact time : 28 day(s)

Degradation : = 0 (±) % after 28 day(s)

Result : under test conditions no biodegradation observed

Deg. product :

Method : other

Year : 1992

GLP : no

Test substance : as prescribed by 1.1 - 1.4

Result : 4,4'-Methylenebis(2-chloroaniline) is not expected to undergo biodegradation based on a screening test. 4,4'-Methylenebis(2-chloroaniline), present at 100 mg/liter, reached 0% of its theoretical BOD in 4 weeks using an activated sludge inoculum at 30 mg/liter and the Japanese MITI test.

Reliability : (2) valid with restrictions
Japanese CITI studies are widely accepted as valid

Flag : Critical study for SIDS endpoint
20.12.2005 (10)

Type : aerobic

Inoculum : activated sludge

Concentration : 2 mg/l related to Test substance related to

Contact time :

Degradation : = 0 (±) % after 42 day(s)

Result :

Deg. product :

Method : other

Year : 1979

GLP : no

Test substance : as prescribed by 1.1 - 1.4

Method : STATIC SYSTEM:
BSA -- Biodegrades slow with acclimation
Rate: 10; 0; 0; 3; 1; 7; 6
Rate Units: % DEGRADATION
Oxygen Condition: AEROBIC
Analysis Method: GC
Incubation Time [Days]: 7
Test Chemical Conc. [ppm]: 2
Acclimation Period [Days]: 7; 14; 21; 28; 35; 42
Dissolved Organic Carbon: YEAST EXTRACT
Microbial Population: 10 (%)
Inoculum: ACTIVATED SLUDGE
Temperature [degrees C]: 25
Remarks: FLASKS SUBCULTURED FROM PREVIOUS WEEK

CONTINUOUS FEED SYSTEM
BFA -- Biodegrades fast with acclimation
Rate: 96; 96; 100; 94; 77; 100
Rate Units: % DEGRADATION
Oxygen Condition: AEROBIC
Continuous/Semi-Continuous Activated Sludge: c
Analysis Method: GC
Retention Time [Days]: 24
Test Chemical Conc. [ppm]: 2
Acclimation Period [Days]: 7; 14; 21; 28; 35; 42
Inoculum: ACTIVATED SLUDGE
Temperature [degrees C]: 25

Result : 4,4'-Methylenebis(2-chloroaniline) was not degraded after 6 weeks in a

3. Environmental Fate and Pathways

Id 101144

Date 22.12.2005

Reliability

20.12.2005

static system in which 2 mg/l of the compound was seeded with sludge, incubated for 7 days, and the process repeated using cultures from the original system. However, when 4,4'-methylenebis(2-chloroaniline) containing water to a continuous feed activated sludge reactor, 96% of the chemical in the water was removed in 24 hours after 1 week of operation.

: (2) valid with restrictions
EPA source document

(18)

4.1 ACUTE/PROLONGED TOXICITY TO FISH

Type :
Species : *Oryzias latipes* (Fish, fresh water)
Exposure period : 96 hour(s)
Unit : mg/l
LC50 : = .61
Method : other
Year : 2001
GLP : no data
Test substance : as prescribed by 1.1 - 1.4

Reliability : (2) valid with restrictions
Japanese MOE studies are widely accepted as valid

Flag : Critical study for SIDS endpoint
21.12.2005

(42)

4.2 ACUTE TOXICITY TO AQUATIC INVERTEBRATES

Type :
Species : *Daphnia magna* (Crustacea)
Exposure period : 48 hour(s)
Unit : mg/l
EC50 : = .92
Method : other
Year : 2001
GLP : no data
Test substance : as prescribed by 1.1 - 1.4

Reliability : (2) valid with restrictions
Japanese MOE studies are widely accepted as valid

Flag : Critical study for SIDS endpoint
21.12.2005

(42)

4.3 TOXICITY TO AQUATIC PLANTS E.G. ALGAE

Species : *Selenastrum capricornutum* (Algae)
Endpoint : other: growth rate and biomass
Exposure period : 72 hour(s)
Unit : mg/l
Method : other
Year : 2001
GLP : no data
Test substance : as prescribed by 1.1 - 1.4

Result : Growth rate:
48 hr ECr50 > 1.9 mg/L; 48 hr NOEC = 1.4

Biomass:
72 hr ECb50 > 1.9 mg/L; 72 hr NOEC = 0.74

Reliability : (2) valid with restrictions
Japanese MOE studies are widely accepted as valid

Flag : Critical study for SIDS endpoint
21.12.2005

(42)

5.1.1 ACUTE ORAL TOXICITY

Type : LD50
 Value : = 1140 mg/kg bw
 Species : rat
 Strain :
 Sex :
 Number of animals :
 Vehicle :
 Doses :
 Method : other
 Year : 1996
 GLP : no data
 Test substance : as prescribed by 1.1 - 1.4

Reliability : (2) valid with restrictions
 Source is peer-reviewed published data

Flag : Critical study for SIDS endpoint
 19.12.2005

(28)

Type : LD0
 Value : = 640 mg/kg bw
 Species : mouse
 Strain :
 Sex :
 Number of animals :
 Vehicle :
 Doses :
 Method : other
 Year : 1996
 GLP : no data
 Test substance : as prescribed by 1.1 - 1.4

Reliability : (2) valid with restrictions
 Source is peer-reviewed published data

19.12.2005

(28)

Type : other
 Value :
 Species : mouse
 Strain :
 Sex :
 Number of animals :
 Vehicle :
 Doses :
 Method : other
 Year : 2000
 GLP : no data
 Test substance : as prescribed by 1.1 - 1.4

Result : In mice, poisoning is accompanied by adynamia, apathy, loss of appetite. Death occurs in 3 days. Manifestations of the toxic action in rats are not very pronounced and death occurs in 2 days. The LOEL for methemoglobinemia formation appears to be 83 mg/kg bw. Gross pathology examination revealed distention of the stomach and intestine, traces of blood in the urinary bladder, and pleural effusion in the thorax. Histology examination detected fine-drop adiposis of the liver, tiny foci of inflammatory infiltration, and circulatory disturbances in the visceral organs.

Reliability : (2) valid with restrictions

20.12.2005

Source is peer-reviewed published data

(49)

5.1.2 ACUTE INHALATION TOXICITY**5.1.3 ACUTE DERMAL TOXICITY**

Type : LD50
Value : > 5000 - mg/kg bw
Species : rabbit
Strain :
Sex :
Number of animals :
Vehicle :
Doses :
Method : other
Year : 1996
GLP : no data
Test substance : as prescribed by 1.1 - 1.4

Reliability : (2) valid with restrictions
Source is peer-reviewed published data

21.12.2005

(28)

5.1.4 ACUTE TOXICITY, OTHER ROUTES**5.4 REPEATED DOSE TOXICITY****5.5 GENETIC TOXICITY 'IN VITRO'**

Type : Bacterial reverse mutation assay
System of testing : Salmonella typhimurium TA98 and TA100; Escherichia coli WP2uvrA
Test concentration :
Cycotoxic concentr. :
Metabolic activation : with
Result : positive
Method : other
Year : 1994
GLP : no data
Test substance : as prescribed by 1.1 - 1.4

Remark : Results excerpted from ATSDR (1994) Toxicological Profile for 4,4'-Methylene-bis(2-chloroaniline) MBOCA. U.S. Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry

Result : In vitro testing has provided clear and convincing evidence that MBOCA is mutagenic in the Salmonella typhimurium/mammalian microsome mutagenesis assay and that the mutagenic effect requires exogenous metabolic activation (Baker and Bonin 1981; Cocker et al. 1985; Dunkel et al. 1984; MacDonald 1981; Martire et al. 1981; McCann et al. 1975; Messerly et al. 1987; Rao et al. 1982). Although not all investigators used each tester strain, the general result is that MBOCA is mutagenic only in strains TA98 and TA100, at 250 µg/plate, with some inconsistency regarding strain TA98. MBOCA and its metabolites are not mutagenic in S.

typhimurium strains TA1535, TA1537, or TA1538. This suggests that the mutagenic effect of MBOCA metabolites in some bacteria is dependent on the plasmid pKM101; strains TA98 and TA100 contain this plasmid, but strains TA1535, TA1537, and TA1538 do not (Ames et al. 1975). This hypothesis is supported by the finding that S9-activated MBOCA is mutagenic in *Escherichia coli* strain WP2uvrA only in the presence of the plasmid pKM101 (Matsushima et al. 1981). The plasmid carries genes involved in an "errorprone" DNA repair system that introduces mutations as it removes DNA damage (Walker 1984). S9 derived from dog and human liver could activate MBOCA to a form mutagenic to strain TA100 but only in a protocol using a fluctuation assay (Cocker et al. 1985).

Most of MBOCA's mutagenic activity appears to be due to the N-hydroxy metabolite, which caused dose-dependent increases in mutations of *S. typhimurium* strains TA100 and TA98 in a pre-incubation assay using nonactivated doses of $\approx 5 \mu\text{g}/\text{plate}$ (Kuslikis et al. 1991). This metabolite is produced by several species, including dogs and humans (Butler et al. 1989; Chen et al. 1989; Morton et al. 1988). The mononitroso derivative appears to be direct-acting mutagen but is much less potent, causing a statistically significant revertant increase in the pre-incubation assay at the highest tested nontoxic dose ($50 \mu\text{g}/\text{plate}$). Neither the o-hydroxy nor the dinitroso derivatives were direct-acting mutagens at up to 50 or 500 $\mu\text{g}/\text{plate}$, respectively. Neither chemical was tested to cytotoxic levels (Kuslikis et al. 1991). N-acetylation is a deactivating step. Neither n-acetyl nor N,N-diacetyl derivatives were mutagenic in *S. typhimurium* in the absence of activation (Hesbert et al. 1985). In the presence of S9 activation, the mutagenic activity of the acetylated derivatives is less than that of the parent compound (Cocker et al. 1986; Hesbert et al. 1985).

Conclusion	: In vitro testing has provided clear and convincing evidence that MBOCA is mutagenic in the <i>Salmonella typhimurium</i> /mammalian microsome mutagenesis assay and that the mutagenic effect requires exogenous metabolic activation
Reliability	: (2) valid with restrictions Source is peer-reviewed published data
Flag 21.12.2005	: Critical study for SIDS endpoint (3) (4) (5) (6) (11) (12) (13) (15) (21) (26) (29) (31) (32) (33) (37) (40) (45) (62)
Type	: DNA damage and repair assay
System of testing	:
Test concentration	:
Cytotoxic concentr.	:
Metabolic activation	:
Result	: positive
Method	: other
Year	: 1994
GLP	: no data
Test substance	: as prescribed by 1.1 - 1.4
Remark	: Results excerpted from ATSDR (1994) Toxicological Profile for 4,4'-Methylene-bis(2-chloroaniline) MBOCA. U.S. Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry
Result	: DNA adducts have been found following oral (Cheever et al. 1990; Kugler-Steigmeier et al. 1989; Segerback and Kadlubar 1992) and dermal (Cheever et al. 1990) administration of radiolabeled MBOCA to rats, following the incubation of radiolabeled MBOCA with explants of dog and human bladder urothelium (transitional cell epithelium) (Stoner et al. 1988) and incubation of rat DNA and radiolabeled N-hydroxy-MBOCA (Segerback and Kadlubar 1992). The level of binding increased with dose, but the increase was not linear. Considerable individual variation in binding levels, varying over at least a 10-fold range, was found in both dogs and humans. At least six adducts were found in dog bladder epithelium; four adducts

	<p>were found in human bladder epithelium, three of which appeared to be the same as those found in dogs. DNA adduct formation in dog bladder tissue is of particular note, since MBOCA has been found to cause bladder tumors in dogs (Stula et al. 1977).</p>
Reliability	: (2) valid with restrictions Source is peer-reviewed published data
Flag	: Critical study for SIDS endpoint
20.12.2005	(4) (8) (25) (48) (52) (53)

5.6 GENETIC TOXICITY 'IN VIVO'

Type	: other
Species	:
Sex	:
Strain	:
Route of admin.	:
Exposure period	:
Doses	:
Result	: positive
Method	: other
Year	: 1994
GLP	: no data
Test substance	: as prescribed by 1.1 - 1.4

Remark	: Results excerpted from ATSDR (1994) Toxicological Profile for 4,4'-Methylene-bis(2-chloroaniline) MBOCA. U.S. Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry
Result	: In vivo animal studies provide direct and indirect evidence that MBOCA is a mutagen; MBOCA metabolites were bound to DNA following oral (Cheever et al. 1990; Kugler-Steigmeier et al. 1989) or dermal (Cheever et al. 1990) exposure in rats. Small increases in the SLRL values were observed in <i>D. melanogaster</i> adults fed a 7.5 millimolar MBOCA solution for 3 days (Vogel et al. 1981).

MBOCA induced gene mutations at the thymidine kinase (TK) locus in mouse lymphoma cells (Caspary et al. 1988; Myhr and Caspary 1988). Unscheduled DNA synthesis (UDS) was induced in HeLa cells (Martin and McDermid 1981), in rat primary hepatocytes at $>10 \mu\text{mol}$ (McQueen et al. 1981; Mori et al. 1988; Williams et al. 1982) and in hamster (McQueen et al. 1981) and rabbit (McQueen and Williams 1987) hepatocytes. The concentration that tested positive in the mouse was $50 \mu\text{mol}$ (McQueen et al. 1981). Sensitivity to MBOCA showed species-specific variations: rat $>$ mouse $>$ hamster $>$ rabbit (McQueen et al. 1981, 1983). Because hepatocytes have their own metabolic activation systems, no exogenous metabolic activation is needed. In assays using attachment independence as an end point, MBOCA, at concentrations near the LC50, transformed baby hamster kidney (Daniel and Dehnel 1981; Styles 1981), rat embryo (Dunkel et al. 1981; Traul et al. 1981), and Balb/3T3 cells (Dunkel et al. 1981).

Transformation assays have not been evaluated as thoroughly as some other genotoxicity assays. In an interlaboratory comparison, one laboratory found equivocal evidence that nonactivated MBOCA induced sister chromatid exchange in Chinese hamster ovary cells at a dose of $5.0 \mu\text{g/mL}$. The response is considered equivocal because the dose-response curve was inconsistent. The result was not confirmed by the second laboratory. In the presence of S9 activation, positive results were obtained by one laboratory at $50 \mu\text{g/mL}$. The other laboratory observed the beginning of a dose-response curve, but the high-dose ($30 \mu\text{g/mL}$) results

did not meet the testing laboratory's criterion for a positive response. In a chromosome aberration assay testing activated and nonactivated concentrations up to 5 and 30 µg/mL, respectively, neither laboratory found a positive result (Galloway et al. 1985). Another study (Perry and Thomson 1981) found no evidence of sister chromatid exchange at up to cytotoxic doses (100 µg/mL), but no doses were tested in the probable sensitive range (>10 and <100 µg/mL). Thus, MBOCA is clastogenic in some systems but not others.

In another study, MBOCA was classified as a clastogen on the basis of results from an in vivo micronucleus bone marrow assay (Katz et al. 1981). Evaluations using a two-phase micronucleus assay and intraperitoneal dosing of B6C3F1 hybrid mice were incomplete, but the results suggested that MBOCA is an in vivo clastogen at doses of >32 mg/kg (Salamone et al. 1981) or 50% of the 1 LD50 (Katz et al. 1981). Results were inconclusive because of toxicity during the first phase of testing in which mice were dosed at 0 and 24 hours with 80% of the LD50 (51 mg/kg), and results were evaluated at 48 hours. In the second phase of testing, mice were dosed with 32 or 51 mg/kg and sampled 48 hours later, or in a separate test, dosed with 32 or 48 mg/kg and sampled 36 hours later. Results were negative at 32 mg/kg when sampled at 48 hours but positive at the same dose in a separate assay when sampled at 36 hours. Results were positive at the higher dose for both sampling times. Multiple sampling times were not performed in any assay (Salamone et al. 1981). In another study, results in the micronucleus test were negative at doses up to 50% of the LD50 (32 mg/kg) using CD-1 mice (dosing was done intraperitoneally at 0 and 24 hours and sampling at 30 hours) (Tsuchimoto and Matter 1981).

Conclusion	:	n vivo animal studies provide direct and indirect evidence that MBOCA is a mutagen.
Reliability	:	(2) valid with restrictions Source is peer-reviewed published data
Flag	:	Critical study for SIDS endpoint
21.12.2005		(4) (7) (8) (14) (15) (19) (23) (25) (30) (34) (35) (36) (39) (41) (44) (47) (55) (57) (58) (60) (63)

5.8.1 TOXICITY TO FERTILITY

5.8.2 DEVELOPMENTAL TOXICITY/TERATOGENICITY

-
- (1) ACGIH (2001) American Conference of Governmental Industrial Hygienists. Documentation of Threshold Limit Values for Chemical Substances and Physical Agents and Biological Exposure Indices for 2001. Cincinnati, OH. 2001. p. 1
 - (2) Aldrich Handbook of Fine Chemicals and Laboratory Equipment (2000-2001), Aldrich Chemical Co., Milwaukee, WI., 2000
 - (3) Ames BN, McCann J, Yamasaki E. (1975) Methods for detecting carcinogens and mutagens with the Salmonella/mammalian-microsome mutagenicity test. *Mutation Res* 31:347-363.
 - (4) ATSDR (1994) Toxicological Profile for 4,4'-Methylene-bis(2-chloroaniline) MBOCA. U.S. Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry
 - (5) Baker RS, Bonin AM. (1981) Study of 42 coded compounds with the Salmonella/mammalian microsome assay. In: deserres FJ, Ashby J, eds. Evaluation of short-term tests for carcinogens: Report of the International Collaborative Program. *Prog Mutat Res* 1:249-260.
 - (6) Butler MA, Guengerich FP, Kadlubar FF. (1989) Metabolic oxidation of the carcinogens 4-aminobiphenyl and 4,4'-methylene-bis(2-chloroaniline) by human hepatic microsomes and by purified rat hepatic cytochrome P-450 monooxygenases. *Cancer Res* 49(1):25-31.
 - (7) Caspary WJ, Daston DS, Myhr BC, et al. (1988) Evaluation of the L5178Y mouse lymphoma cell mutagenesis assay: Interlaboratory reproducibility and assessment. *Environmental Molecular Mutagenesis* 12 (suppl 13): 195-229.
 - (8) Cheever KL, Richards DE, Weigel WW, et al. (1990) 4,4'-Methylene-bis(2-chloroaniline) (MOCA): Comparison of macromolecular adduct formation after oral or dermal administration in the rat. *Fundam Appl Toxicol* 14(2):273-28X
 - (10) Chemicals Inspection and Testing Institute (1992) Biodegradation and bioaccumulation data of existing chemicals based on the CSCL Japan. Japan Chemical Industry Ecology - Toxicology and Information Center. ISBN 4-89074-101-1
 - (11) Chen TH, Kuslikis BI, Braselton WE Jr. (1989) Hydroxylation of 4,4'-methylenebis(2-chloroaniline) by canine, guinea pig, and rat liver microsomes. *Drug Metab Dispos* 17(4):406-413.
 - (12) Cocker J, Boobis AR, Davis DS. (1986) Routes of activation of 4,4'-methylenebis(2-chloroaniline) and 4,4'-methylenedianiline to bacterial mutagens. In: International Conference on Practical in vitro Toxicology, Berkshire, England, September 18-20, 1985. *Food Chem Toxicol* 24(6-7):755-756.
 - (13) Cocker J, Boobis AR, Gibson JF, et al. (1985) The metabolic activation of 4,4'-methylenebis(2-chlorobenzeneamine) to a bacterial mutagen by hepatic postmitochondrial supernatant from human and other species. *Environ Mutagen* 7(4):501-509.
 - (14) Daniel MR, Dehnel JM. (1981) Cell transformation test with baby hamster kidney cells. In: deserres FJ, Ashby J, eds. Evaluation of short-term tests for carcinogens: Report of the International Collaborative Program. *Prog Mutat Res* 1:626-637.
 - (15) Dunkel VC, Zeiger E, Brusick D, et al. (1984) Reproducibility of microbial mutagenicity assays: 1. Tests with Salmonella typhimurium and Escherichia coli using a standardized protocol. *Environ mutagen* G(suppl 2):1-254.
 - (16) Ellington, J.J. (1989) Hydrolysis Rate Constants for Enhancing Property - Reactivity Relationships. Report 1989, USEPA/600/3-89/063, NTIS PB89-220479

-
- (17) EPA. (1988) Measurement of hydrolysis rate constants for evaluation of hazardous waste land disposal. Volume 3: Data on 70 chemicals. Athens, GA: U.S. Environmental Protection Agency, Office of Research and Development. EPA/600/3-88028, document no. PB88-234042
- (18) Fochtman, E.G. and Eisenberg, W. (1979) Treatability of carcinogenic and other hazardous organic compounds p.61 USEPA-600/2-79-097
- (19) Galloway SM, Bloom AD, Resnick M, et al. (1985) Development of a standard protocol for in vitro cytogenetic testing with Chinese hamster ovary cells: Comparison of results for 22 compounds in two laboratories. *Environ Mutagen* 21-51.
- (21) Hesbert A, Bottin MC, DeCeaurrez J. (1985) Mutagenicity of 4,4'-methylenebis(2-chloroaniline) "MOCA" and its n-acetyl derivatives in *S. yphimurium*. *Int Arch Occup Environ Health* 55(2):169-174.
- (23) Katz M, Heddle JA, Salamone MF. (1981) Mutagenic activity of polycyclic aromatic hydrocarbons and other Environmental pollutants. In: Dennis AJ, ed. Chemical analytical and biological fate: Polynuclear aromatic hydrocarbons, 5th International Symp. Columbus, OH: Battelle Press, 519-528.
- (25) Kugler-Stegmeier ME, Friederich U, Graf U, et al. (1989) Genotoxicity of aniline derivatives in various short-term tests. *Mutat Res* 211(2):279-289.
- (26) Kuslikis BI, Tosko JE, Braselton WE Jr. (1991) Mutagenicity and effect on gap-junctional intercellular communication of 4,4'-methylenebis(2-chloroaniline) and its oxidized metabolites. *Mutagenesis* 6(1):19-24.
- (27) Leo, A.J. (1978) Title: Report on the Calculation of Octanol/Water Log P Values for Structures in EPA Files
- (28) Lewis, R.J. (1996) Sax's Dangerous Properties of Industrial Materials. 9th ed. Volumes 1-3. New York, NY: Van Nostrand Reinhold, 1996., p. 2228
- (29) MacDonald DJ. (1981) Salmonella/microsome tests on 42 coded chemicals. In: deSerres FJ, Ashby J, eds. Evaluation of short-term tests for carcinogens: Report of the International Collaborative Program. *Prog Mutat Res* 1:285-297.
- (30) Martin CN, McDermid AC. (1981) Testing of 42 coded compounds for their ability to induce unscheduled DNA repair synthesis in HeLa cells. In: deSerres FJ, Ashby J, eds. Evaluation of short-term tests for carcinogens: Report of the International Collaborative Program. *Prog Mutat Res* 1:533-537.
- (31) Martire G, Vricella G, Perfumo Am, et al. (1981) Evaluation of the mutagenic activity of coded compounds in the Salmonella test. In: deSerres FJ, Ashby J, eds. Evaluation of short-term tests for carcinogens: Report of the International Collaborative Program. *Prog Mutat Res* 1:271-279.
- (32) Matsushima T, Takamoto Y, Shirai A, et al. (1981) Reverse mutation test on 42 coded compounds with the *E. coli* WP2 system. In: deSerres FJ, Ashby J, eds. Evaluation of short-term tests for carcinogens: Report of the International Collaborative Program. *Prog Mutat Res* 1:387-395.
- (33) McCann J, Choi E, Yamasaki E, et al. (1975) Detection of carcinogens as mutagens in the Salmonella/microsome test: Assay of 300 chemicals. *Proc natl Acad Sci* 72:5135-5139.
- (34) McQueen CA, MasLansky CJ, Crescenzi SB, et al. (1981) The genotoxicity of 4,4'-methylenebis-2-chloroaniline in rat, mouse and hamster hepatocytes. *Toxicol Appl Pharmacol* 58(2):231-235.

-
- (35) McQueen CA, Maslansky CJ, Williams GM. (1983) Role of the acetylation polymorphism in determining susceptibility of cultured rabbit hepatocytes to DNA damage by aromatic amines. *Cancer Res* 43(7): 3120-3123.
- (36) McQueen CA, Williams GM. (1987) The hepatocyte primary culture/DNA repair test using hepatocytes from several species. *Cell Biology and Toxicology* 3(2):209-218.
- (37) Messerly EA, Fekete JE, Wade DR, et al. (1987) Structure-mutagenicity relationships of benzidine analogues. *Environ Mol Mutagen* 10(3):263-274.
- (38) Meylan, W.M. and Howard, P.H. (1993) *Chemosphere* 26: 2293-99.
- (39) Mori H, Yoshimi n, Sugie S, et al. (1988) Genotoxicity of epoxy resin hardeners in the hepatocyte primary culture/DNA repair test. *Mutat Res* 204(4):683-688.
- (40) Morton KC, Lee M-S, Siedlik P, et al. (1988) Metabolism of 4,4'-methylenebis-2-chloroaniline (MOCA) by rats in Vivo and formation of n-hydroxy MOCA by rat and human liver microsomes. *carcinogenesis* 9(5):731-739.
- (41) Myhr BC, Caspary WJ. (1988) Evaluation of the L5178Y mouse lymphoma cell mutagenesis assay: Intralaboratory results for sixty-three coded chemicals tested at Litton Bionetics, Inc. *Environ Mol Mutagen* 12(Suppl 13):103-194.
- (42) National Institute of Technology and Evaluation, Ministry of the Environment (2001)
- (43) NRC. (1981) Aromatic Amines: An assessment of the biological and Environmental effects. Washington, DC: National Academy Press, p. 19, 168-197.
- (44) Perry PE, Thomson EJ. (1981) Evaluation of the sister chromatid exchange method in mammalian cells as a screening system for carcinogens. In: deSerres FJ, Ashby J, eds. *Evaluation of short-term tests for carcinogens: Report of the International Collaborative Program*. *Prog Mutat Res* 1:560-569.
- (45) Rao TK, Dorsey GF, Allen BE, et al. (1982) Mutagenicity of 4,4'-methylenedianiline derivatives in the Salmonella histidine reversion assay. *Arch Toxicol* 49(3-4):185-190.
- (47) Salamone MF, Heddle JA, Katz M. (1981) Mutagenic activity of 41 compounds in the in vivo micronucleus assay. In: deSerres FJ, Ashby J, eds. *Evaluation of short-term tests for carcinogens: Report of the International Collaborative Program*. *Prog Mutat Res* 1:686-697.
- (48) Segerback D, Kadlubar FF. (1992) Characterization of 4,4'-methylenebis(2-chloroaniline)-DNA adducts formed in vivo and in vitro. *Carcinogenesis* 13:1587-1592.
- (49) Sheftel, V.O. (2000) Indirect Food Additives and Polymers. Migration and Toxicology. Lewis Publishers, Boca Raton, FL. p. 489.
- (50) Smith AM, Woodward Kn. (1983) 4,4'-Methylene bis-(2-chloroaniline)(MBOCA). *Toxic Rev* 8, Part 2:1-19.
- (51) SRC (1988) Syracuse Research Corporation Calculated Values, Reference Number 255435;
- (52) Stoner GD, Shivapurkar nM, Schut HAJ, et al. (1988) DNA binding and adduct formation of 4,4'-methylene-bis(2-chloroaniline) (MOCA) in explant cultures of human and dog bladder. In: King CM, Roman0 LJ, and Schultze D, eds. *Carcinogenic and mutagenic responses to aromatic amines and nitroarines*. New York, NY: Elsevier, 237-240.
- (53) Stula EF, Barnes JR, Sherman H, et al. (1977) Urinary bladder tumors in dogs from 4,4'-methylenebis(2-chloroaniline) (MOCA). *J Environ Pathol Toxicol* 1(1):3 1-50.

-
- (55) Styles JA. (1981) Activity of 42 coded compounds in the BHK-21 cell transformation test. In: deserre FJ, Ashby J, eds. Evaluation of short-term tests for carcinogens: Report of the International Collaborative Program. Prog Mutat Res 1:638-646.
- (56) The Merck Index (1983) An Encyclopedia of Chemicals and Drugs 10th Ed. Windholz, M. Ed. Rahway, N.J.: Merck and Co., Inc.
- (57) Traul KA, Takayama K, Kachevsky V, et al. (1981) A rapid in vitro assay for carcinogenicity of chemical substances in mammalian cells utilizing an attachment-independence endpoint: 2. Assay validation. J Appl Toxicol 1(3):190-195.
- (58) Tsuchimoto T, Matter BE. (1981) Activity of coded compounds in the micronucleus test. In: deserre FJ, Ashby J, eds. Evaluation of short-term tests for carcinogens: Report of the International Collaborative Program. Prog Mutat Res 1:705-711.
- (59) US EPA (2003) Estimation Program Interface (EPI) Suite. Ver.3.11. June 10, 2003. Available at <http://www.epa.gov/oppt/exposure/docs/episuitedi.htm> as of Jan, 2004
- (60) Vogel E, Blijleven WG, Kortselius MJ, et al. (1981) Mutagenic activity of 17 coded compounds in the sex-linked recessive lethal test in *Drosophila melanogaster*. In: deserre FJ, Ashby J. eds. Evaluation of short-term tests for carcinogens: Report of the International Collaborative Program. Prog Mutat Res 1:660-665.
- (61) Voorman, R. and Penner, D. (1986) Fate of MBOCA (4,4'-Methylene-bis(2-chloroaniline)) in soil. Archives of Environmental Contamination and Toxicology, 15: 595-602
- (62) Walker CG. (1984) Mutagenesis and inducible responses to deoxyribonucleic acid damage in *Escherichia coli*. Microbial Rev 48(1):60-9X
- (63) Williams GM, Laspia MF, Dunkel VC. (1982) Reliability of the hepatocyte primary culture/DNA repair test in testing of coded carcinogens and noncarcinogens. Mutat Res 97(5):359-370.

APPENDIX 2
EPIWIN RESULTS

SMILES : Nc(c(cc(c1)Cc(ccc(N)c2CL)c2)CL)c1
CHEM : Benzenamine, 4,4'-methylenebis 2-chloro-
CAS NUM: 000101-14-4
MOL FOR: C13 H12 CL2 N2
MOL WT : 267.16

----- EPI SUMMARY (v3.11) -----

Physical Property Inputs:

Water Solubility (mg/L): -----
Vapor Pressure (mm Hg) : -----
Henry LC (atm-m3/mole) : -----
Log Kow (octanol-water): -----
Boiling Point (deg C) : -----
Melting Point (deg C) : -----

Log Octanol-Water Partition Coef (SRC):

Log Kow (KOWWIN v1.67 estimate) = 3.47
Log Kow (Exper. database match) = 3.91
Exper. Ref: Chem Inspect Test Inst (1992)

Boiling Pt, Melting Pt, Vapor Pressure Estimations (MPBPWIN v1.41):

Boiling Pt (deg C): 404.79 (Adapted Stein & Brown method)
Melting Pt (deg C): 156.63 (Mean or Weighted MP)
VP(mm Hg,25 deg C): 3.93E-006 (Modified Grain method)
MP (exp database): 110 deg C
BP (exp database): 378.9 deg C

Water Solubility Estimate from Log Kow (WSKOW v1.41):

Water Solubility at 25 deg C (mg/L): 8.684
log Kow used: 3.91 (expkow database)
no-melting pt equation used
Water Sol (Exper. database match) = 13.9 mg/L (24 deg C)
Exper. Ref: VOORMAN,R & PENNER,D (1986A)

Water Sol Estimate from Fragments:

Wat Sol (v1.01 est) = 4.5958 mg/L
Wat Sol (Exper. database match) = 13.90
Exper. Ref: VOORMAN,R & PENNER,D (1986A)

ECOSAR Class Program (ECOSAR v0.99g):

Class(es) found:
Aromatic Amines

Henrys Law Constant (25 deg C) [HENRYWIN v3.10]:

Bond Method : 3.29E-011 atm-m3/mole
Group Method: 1.14E-011 atm-m3/mole
Henrys LC [VP/WSol estimate using EPI values]: 1.591E-007 atm-m3/mole

Probability of Rapid Biodegradation (BIOWIN v4.01):

Linear Model : -0.1573
Non-Linear Model : 0.0003

Expert Survey Biodegradation Results:

Ultimate Survey Model: 1.8508 (months)
Primary Survey Model : 2.8463 (weeks)

Readily Biodegradable Probability (MITI Model):

Linear Model : -0.3921
Non-Linear Model : 0.0006

Atmospheric Oxidation (25 deg C) [AopWin v1.91]:

Hydroxyl Radicals Reaction:

OVERALL OH Rate Constant = 77.5166 E-12 cm3/molecule-sec

Half-Life = 0.138 Days (12-hr day; 1.5E6 OH/cm3)

Half-Life = 1.656 Hrs

Ozone Reaction:

No Ozone Reaction Estimation

Soil Adsorption Coefficient (PCKOCWIN v1.66):

Koc : 1.353E+004

Log Koc: 4.131

Aqueous Base/Acid-Catalyzed Hydrolysis (25 deg C) [HYDROWIN v1.67]:

Rate constants can NOT be estimated for this structure!

BCF Estimate from Log Kow (BCFWIN v2.15):

Log BCF = 2.311 (BCF = 204.5)

log Kow used: 3.91 (expkow database)

Volatilization from Water:

Henry LC: 1.14E-011 atm-m3/mole (estimated by Group SAR Method)

Half-Life from Model River: 8.394E+007 hours (3.498E+006 days)

Half-Life from Model Lake : 9.158E+008 hours (3.816E+007 days)

Removal In Wastewater Treatment (recommended maximum 99%):

Total removal: 26.13 percent

Total biodegradation: 0.29 percent

Total sludge adsorption: 25.84 percent

Total to Air: 0.00 percent

Level III Fugacity Model:

	Mass Amount (percent)	Half-Life (hr)	Emissions (kg/hr)
Air	9.87e-005	3.31	1000
Water	15.5	1.44e+003	1000
Soil	82.1	1.44e+003	1000
Sediment	2.42	5.76e+003	0

Persistence Time: 1.59e+003 hr
